

Mental health considerations for older adults and individuals receiving care in facility-based settings during the COVID-19 pandemic

Updated 04/22/2020



The information below has been selected to support providers with patients in facility-based care settings who may be experiencing increased symptoms of anxiety, depression and suicidality, particularly due to circumstances related to COVID-19 precautions.

General health guidance

Older adults and people with severe underlying medical conditions are at higher risk for developing more serious complications from COVID-19 illness. Read more about the NIH updates on the virus and federal response: [nih.gov/health-information/coronavirus](https://www.nih.gov/health-information/coronavirus)

The CDC has issued guidance on how to best protect older adults during the pandemic, including recommendations to stay home, wash hands frequently and maintain a physical distance of six feet or more from others in public spaces: [cdc.gov/coronavirus/2019-ncov/index.html](https://www.cdc.gov/coronavirus/2019-ncov/index.html)

Emotional impact on older adults

The outbreak of COVID-19 may be emotionally stressful for older adults. Stressors for older adults during an infectious disease outbreak can include:

- Physical health, pain and disability.
- Social isolation, feeling lonely or disconnected. This may be increased for those in skilled nursing facilities and/or long-term care homes.
- Disruption in routine, including changes in daily life, inability to connect with family and difficulty with grocery shopping.
- Losses.
- Mental health conditions.

Signs of stress anxiety, depression or suicidality in older adults may include:

- Increased fear and worry about their health and the health of their loved ones.
- Changes in sleep or eating patterns.
- Difficulty sleeping or concentrating.
- Worsening of chronic health problems.
- Worsening of mental health conditions.
- Self-isolating.

How can you help?

The American Foundation for Suicide Prevention has issued tips for caregivers, health care workers and loved ones to support elders during the pandemic:

- Empathize and validate worries.
- Support connections with loved ones.
- Protect yourself and others from illness.
- Encourage engagement in old/new activities.
- Know your local resources and supports.
- Consider administering a validated suicide risk assessment.
- Familiarize yourself with your organization's workflow for suspected depression, anxiety or other mental health concerns.

Local mental health resources for older adults:

- **Oregon Psychiatric Access Line** | ohsu.edu/school-of-medicine/child-and-adolescent-psychiatry/oregon-psychiatric-access-line

Free, same-day psychiatric consultation service for clinical teams through a call center operated Monday-Friday, 9 a.m. to 5 p.m. The toll-free number is 855-966-7255.

- **Senior Loneliness Line** | seniorlonelinessline.org/

A free, confidential warmline staffed 24/7 for Oregonians aged 55 and older who are experiencing loneliness, isolation, depression or anxiety. The statewide number is 503-200-1633.

Validated suicide risk screening tools:

| Tool | Link | Description |
|--|---|---|
| Ask Suicide-Screening Questions (ASQ) | nimh.nih.gov/research/research-conducted-at-nimh/asq-toolkit-materials/asq-tool/information-sheet_155866.pdf | A four-item questionnaire designed for administration by non-clinicians in physical health settings for all ages. |
| Patient Safety Screener-3 (PSS-3) | sprc.org/microlearning/patientsafetyscreener | A three-item questionnaire validated for administration in emergency departments by all staff, for all adult populations. |
| Columbia Suicide Severity Scale (C-SSRS) | http://cssrs.columbia.edu/ | An extensive, two-page questionnaire intended for use by multi-disciplinary staff across education levels to inventory current and past suicidality and predictive risk factors. Endorsed by CDC, NIH, FDA, SAMHSA. |

National mental health resources for older adults

Suicide prevention:

- **[Issue Brief 4: Preventing Suicide in Older Adults](#)**
Source: National Council on Aging (NCOA)
Description: List of risk factors for completed suicide and prevention strategies to be taken by aging services workers, behavioral health providers and primary care providers.
- **[Promoting Emotional Health and Preventing Suicide: A Toolkit for Senior Centers](#)**
Source: Substance Abuse and Mental Health Services Administration (SAMHSA)
Description: Guide to early warning signs and early intervention of suicidal ideation in community senior centers.
- **[Promoting Emotional Health and Preventing Suicide: A Toolkit for Senior Living Communities](#)**
Source: SAMHSA
Description: Guide to prevention and early intervention of suicidal ideation across the continuum of senior living communities.
- **[Weaving a Community Safety Net to Prevent Older Adult Suicide](#)**
Source: NASMHPD
Description: Comprehensive article of best practices for elder-specific suicide risk assessment and prevention across healthcare and community settings.