



Farxiga (dapagliflozin)
 Requires PA for CHF, CKD stage 2-4, DM with ASCVD

- [Covered DM Supplies](#)
- [OHP PA Criteria](#)
- [OHP Formulary](#)
- [Medicare Diabetes Coverage](#)

Formulary Status	Max day supply	Class	Medications	Dosing	Considerations
Formulary	90	Biguanide	Metformin IR and ER	500-2g total daily dose	Preferred initial therapy for most DM2 patients
Formulary	90	TZD	Pioglitazone	15-45 mg once daily	Can cause or exacerbate CHF in some patients – avoid use in those patients
Formulary	90	Sulfonylurea	Glimepiride, Glipizide, Glyburide	Varies	Can cause hypoglycemia
Step Therapy	34	SGLT-2 Inhibitors	Steglatro	5-15mg once daily	Dapagliflozin available with PA Steglatro Step Criteria Dapagliflozin PA Criteria
Step Therapy	34	DPP-4 inhibitors	Alogliptin	25mg once daily	Alogliptin Step Criteria
PA	28	GLP-1 Agonists	Ozempic, Victoza, Monjaro, Trulicity, Byetta, Bydureon	See FDA label for DM treatment doses	GLP-1 PA Criteria
Insulin					
Formulary	90	Long-acting/intermediate acting insulin	Lantus, Glargine YFGN, Rezvoglar, Novolin N, Humulin N	Varies	Pen and vials covered when applicable
Formulary	90	Short-acting insulin	Admelog, Insulin Lispro, Novolin R, Humulin R	Varies	Pen and vials covered when applicable
PA	90	Short-acting insulin	Insulin Lispro Junior Pen Injector	Varies	For half-unit dosage, generic is preferred product, requiring PA.

- Statins are generally recommended for patients over 40 years old with diabetes to reduce the risk of ASCVD.
- Diet support: CareOregon has paired with Foodsmart to provide our members with free dietary support including meal planning with a dietician.

More info can be found here: <https://www.careoregon.org/members/more-careoregon-services/foodsmart-nutrition-program>